

ANIL P. PUNJABI, M.D., D.D.S.
Board Certified
American Board of Plastic Surgery
American Board of Oral and Maxillofacial Surgery
365 Terracina Blvd., Redlands, CA 92373

POST OPERATIVE INSTRUCTIONS FOR INJECTABLE FILLERS

COLLAGEN/FAT INJECTIONS

INSTRUCTIONS

- Have someone drive you home after surgery and help you at home for 1-2 days.
- Get plenty of rest.
- Follow balanced diet.
- Decreased activity may promote constipation, so you may want to add more raw fruit to your diet, and be sure to increase fluid intake.
- Take pain medication as prescribed. Do not take aspirin or any products containing aspirin.
- Do not drink alcohol when taking pain medications.
- Even when not taking pain medications, no alcohol for 3 weeks as it causes fluid retention.
- If you are taking vitamins with iron, resume these as tolerated.
- Do not smoke, as smoking delays healing and increases the risk of complications.

ACTIVITIES

- Do not drive until you are no longer taking any pain medications (narcotics).
- Start walking as soon as possible, as this helps to reduce swelling and lowers the chance of blood clots.
- Unless stated on this form, discuss your time off work with your surgeon.

TREATED AREA CARE

- You may shower after your first post-op appt.
- Avoid exposing scars to sun for at least 12 months.
- Always use a strong sunblock, if sun exposure is unavoidable (SPF 30 or greater).
- Keep steri-strips on; replace if they come off.
- Inspect daily for signs of infection.
- No tub soaking while sutures or drains are in place.
- You may wear makeup with sunblock protection shortly.
- Stay out of the sun until redness and bruising subsides (usually 48 hours).

WHAT TO EXPECT

- Temporary stinging, throbbing, burning sensation, redness, swelling, bruising, and excess fullness.

Fat:

- Some swelling, bruising or redness in the donor and recipient sites.
- Swelling and puffiness may last several weeks.
- Redness and bruising usually lasts about 48 hours.

Collagen:

- Any redness you may have in the injected site usually disappears within 24 hours; however, in some patients, particularly fair-skinned, it may persist for a week or more.
- Temporary swelling and redness in the treated area should dissipate within a few days.
- Tiny scabs over the needle-stick areas; they generally heal quickly.

APPEARANCE

- Your new appearance may be short-lived (a few weeks) or long lasting. This is natural. Everyone's "staying power" varies.
- Improved skin texture.
- Firmer and smoother skin.

FOLLOW-UP CARE

- With regular follow-up treatments, you can easily maintain your new look.
- Repeated treatments may be necessary.

WHEN TO CALL

- If you have increased swelling or bruising.
- If swelling and redness persist after a few days.
- If you have increased redness along the incision.
- If you have severe or increased pain not relieved by medication.
- If you have any side effects to medications; such as, rash, nausea, headache, vomiting.
- If you have an oral temperature over 100.4 degrees.
- If you have any yellowish or greenish drainage from the incisions or notice a foul odor.
- If you have bleeding from the incisions that is difficult to control with light pressure.
- If you have loss of feeling or motion.
- If you have any sign of abscesses, open sores, skin peeling or lumpiness.

For medical questions, please call:

- **909-798-9950**, Monday – Friday, 8 a.m. – 5 p.m.
- After hours and on weekends for emergency calls – Dr. Punjabi @ **909-213-6183**
on call.