# ANIL P. PUNJABI, M.D., D.D.S.

# **Board Certified**

# American Board of Plastic Surgery American Board of Oral and Maxillofacial Surgery

365 Terracina Blvd., Redlands, CA 92373

#### POST OPERATIVE INSTRUCTIONS FOR FOREHEAD LIFT

### (CLASSIC LIFT; ENDOSCOPIC LIFT)

#### **INSTRUCTIONS**

- Have someone drive you home after surgery and help you at home for 1-2 days.
- Get plenty of rest.
- Plan on taking it easy for at least the first week.
- Follow balanced diet.
- Decreased activity may promote constipation, so you may want to add more raw fruit to your diet, and be sure to increase fluid intake.
- Take pain medication as prescribed. Do not take aspirin or any products containing aspirin.
- Do not drink alcohol when taking pain medications.
- Even when not taking pain medications, no alcohol for 3 weeks as it causes fluid retention.
- If you are taking vitamins with iron, resume these as tolerated.
- Do not smoke, as smoking delays healing and increases the risk of complications.

#### **ACTIVITIES**

- Start walking as soon as possible, this helps to reduce swelling and lowers the chance of blood clots.
- Do not drive until you are no longer taking any pain medications (narcotics).
- Back to work or school in 7-10 days, usually sooner for endoscopic patients.
- Limit vigorous physical activity for several weeks; i.e., heavy housework, sex, bending or any activity that increases your blood pressure.
- Full social activities can usually be resumed in 2 weeks.

#### INCISION CARE

- You may shower after your first post-op appt.
- Avoid exposing scars to sun for at least 12 months.
- Limit exposure to heat or sun for several months.
- Always use a strong sunblock, if sun exposure is unavoidable (SPF 30 or greater).

- Keep steri-strips on; replace if they come off.
- Keep incisions clean and inspect daily for signs of infection.
- Shower and shampoo in 2 days or as soon as bandage is removed.
- When using hair dryer, never set temperature on "hot", as you may not have feeling in operated area.
- Hair bleach or permanent can be applied after complete healing, usually 3-4 weeks.
- Facial makeup can be applied to cover up bruising 3 days after surgery, but not on the incisions.

## Classic Lift:

• Keep your head elevated for 2-3 days after surgery to help keep the swelling down.

#### WHAT TO EXPECT

- May feel tired and let down at first.
- Temporary discomfort, tightness of skin, swelling, bruising, numbness and headaches.
- Full recovery from bruising in 2-3 weeks.

## Classic Lift:

- Swelling may affect cheeks and eyes for a week or so.
- As nerves heal, itching may replace numbness on top of scalp -- May last up to 6 months.
- Possible hair loss.

### Endoscopic Lift:

- Minimal postoperative pain.
- As nerves heal, itching may replace numbness on top of scalp, but there is less of an itching sensation with the endoscopic forehead lift.

#### **APPEARANCE**

- Younger, more rested look.
- Final results realized in approximately 6 months.
- Visible signs of surgery should fade completely in approximately 3 weeks.

## Classic Lift:

 Normal growth of hair around incision usually resumes in a few weeks or months.

#### **FOLLOW-UP CARE**

### Classic Lift:

- Most stitches or clips are removed within 14 days, sometimes in 2 stages.
- If bandages were used, they will be removed in 1-2 days.

# Endoscopic Lift:

- Stitches or staples used to close the incisions are usually removed within a
  week
- Temporary fixation screws are usually removed within 2 weeks.

#### WHEN TO CALL

- If you have increased swelling or bruising.
- If swelling and redness persist after a few days.
- If you have increased redness along the incision.
- If you have severe or increased pain not relieved by medication.
- If you have any side effects to medications; such as, rash, nausea, headache, vomiting.
- If you have an oral temperature over 100.4 degrees.
- If you have any yellowish or greenish drainage from the incisions or notice a foul odor.
- If you have bleeding from the incisions that is difficult to control with light pressure.
- If you have loss of feeling or motion.

# For medical questions, please call:

- **909-798-9950**, Monday Friday, 8 a.m. 5 p.m.
- After hours and on weekends for emergency calls Dr. Punjabi @ 909-213-6183